

JOURNAL OF FUNCTIONAL ANALYSIS 14, 146-161 (1973)

On the Stochastic Maximum Principle in Banach Space

HUI-HSIUNG KUO*

*Department of Mathematics, University of Virginia,
Charlottesville, Virginia 22903*

Communicated by J. L. Lions

Received January 30, 1973

1. INTRODUCTION

We initiate the study of stochastic control theory in infinite dimensional Banach space by proving an infinite dimensional analog of the celebrated Pontryagin's maximum principle [12] and its stochastic version due to Kushner [11]. We prove also an existence theorem for optimal controls. For the finite dimensional theory, see [1-5, 13] and the references in the comprehensive review paper [4]. Our approach in proving the maximum principle is strongly influenced by [13] where the stochastic integral is McShane's belated integral. However, we use a new machinery introduced in [10]. Also, our existence theorem is new even if the Banach space is finite dimensional.

It is well-known that a real separable Banach space can be regarded as an abstract Wiener space [6]. Thus we will work on stochastic systems in a fixed abstract Wiener space $H \subset B$. Consider the following stochastic integral equation

$$X_t = x + \int_0^t A(s, X_s) dW_s + \int_0^t \sigma(s, X_s, u(s)) ds, \quad 0 \leq t \leq \tau, \quad (1)$$

where τ is a fixed time, W_t is a Wiener process in $H \subset B$ and the control function u takes values in a subset U of a separable Banach space G (finite or infinite dimensional). Similar type of Eq. (1) has been studied in [8]. Let

$$C(u, X) = \int_0^\tau f(t, X_t, u(t)) dt + p(X_\tau), \quad (2)$$

* Research supported in part by NSF grant GU-3784.

and

$$\Phi(u, X) = E[C(u, X)] \quad (\text{the expectation}). \quad (3)$$

The regularity assumptions on the functions A , σ , f and p will be made precise in Section 2. Our object is to seek a necessary condition for u_0 such that u_0 and the corresponding trajectory X_0 will minimize Φ over the set of all u and X satisfying Eq. (1). This necessary condition is given in Theorem 1 (Section 2). As in the finite dimensional case it involves the Lagrange multiplier ϕ_t . However, a dissimilarity occurs in infinite dimensional case, namely, ϕ_t is a stochastic process satisfying a linear integral equation in the Hilbert space H (instead of B). Section 3 is devoted to the proof of Theorem 1 and in Section 4 we give two examples, one in which B is infinite dimensional and another B is one-dimensional.

Next, consider the control system (1)–(2) with the diffusion coefficients and the cost function given as follows:

$$\begin{aligned} A(t, x) &= J + K(t, x), & \sigma(t, x, u) &= C(t)x + \theta(t, u), \\ f(t, x, u) &= |Dx|^2 + \langle F(t)x, \zeta(t, u) \rangle + g(t, u), & p &= 0. \end{aligned} \quad (4)$$

Under a rather mild assumption on (4), we have, in Theorem 2 (Section 2), the existence of an optimal control for Φ if we restrict Φ to a certain class of control functions. In fact, this restriction is motivated by Theorem 2.3 of [1]. The proof of Theorem 2 is given in Section 5.

2. MAXIMUM PRINCIPLE AND EXISTENCE THEOREM

Notation: $|\cdot| = H$ -norm, $\|\cdot\| = B$ -norm, $|\cdot|_0 = G$ -norm, $\langle \cdot, \cdot \rangle = H$ -inner product, $(\cdot, \cdot) =$ natural pairing of B^* and B , ($B^* \subset H \subset B$ as in [8]).

The appropriate smoothness for functions defined in the abstract Wiener space $H \subset B$ is the so-called Fréchet differentiability in H -directions (shortly, H -differentiability). A function f from B into another Banach space Z is said to be H -differentiable at x if there exists a (unique) linear operator $T \in L(H, Z)$ (\equiv the Banach space of linear operators from H into Z with operator norm $\|\cdot\|_{H,Z}$) such that $\|f(x+h) - f(x) - Th\|_Z = o(|h|)$, $h \in H$. T will be denoted by $f'(x)$ or f_x (specially, when f depends also on other variables). f is said to be C_H^1 if $f'(x)$ exists for all x in B and f' is continuous from B into $L(H, Z)$.

We now state the hypotheses of A , σ , f and p in Eqs. (1) and (2).

(A-1). $A(t, x) = J + K(t, x)$, where $J \in L(B, B)$ and K is a continuous map from $[0, \tau] \times B$ into $L_2(H)$ (\equiv the Hilbert space of Schmidt operators of H with Schmidt norm $\|\cdot\|_2$).

(A-2). There is a constant c such that $\|K(t, x) - K(t, y)\|_2 \leq c\|x - y\|$, $\|K(t, x)\|_2 \leq c(1 + \|x\|)$ for all t and x .

(A-3). K is C_H^1 in x variable such that K_x is bounded and continuous from $[0, \tau] \times B$ into $L(H, L_2(H))$.

(σ -1). σ is a continuous map from $[0, \tau] \times B \times U$ into H such that $|\sigma(t, x, u) - \sigma(t, y, v)| \leq c(\|x - y\| + \|u - v\|_0)$ and $|\sigma(t, x, u)| \leq c(1 + \|x\| + \|u\|_0)$.

(σ -2). σ is C_H^1 in x variable such that σ_x is bounded and continuous from $[0, \tau] \times B \times U$ into $L(H, H)$.

(f-1). f is a real-valued continuous function in $[0, \tau] \times B \times U$ such that $|f(t, x, u)| \leq c(1 + \|x\|^2 + \|u\|_0^2)$.

(f-2). f is C_H^1 in x variable such that $|f_x(t, x, u)| \leq c(1 + \|x\| + \|u\|_0^2)$.

(p-1). p is a real-valued C_H^1 continuous function in B such that $|p'(x)| \leq c(1 + \|x\|)$.

We now specify the control functions space \mathcal{U} . \mathcal{U} consists of all square integrable functions $u(t)$ in $[0, \tau]$ taking values in U , i.e., $\|u\|_0^2 = \int_0^\tau \|u(t)\|_0^2 dt < \infty$.

The adjoint system of (1)–(2) is defined to be the following linear stochastic differential equation with state space the Hilbert space H ,

$$d\phi_t = -A_x(t, X_t)^* \phi_t dW_t - [\sigma_x(t, X_t, u(t))^* \phi_t + f_x(t, X_t, u(t))] dt \quad (5)$$

$$\phi_\tau = p'(X_\tau),$$

where if $T \in L(H, H)$ then T^* denotes the adjoint operator of T , and if $S \in L(H, L(H, H))$ then $S^* \in L(H, L(H, H))$ is defined as follows: $S^*(h) = [S(h)]^*$, $h \in H$. The Hamiltonian \mathcal{H} is defined for each t in $[0, \tau]$ to be a function in $H \times B \times U$ as follows:

$$\mathcal{H}(\phi, x, u; t) = f(t, x, u) + \langle \phi, \sigma(t, x, u) \rangle. \quad (6)$$

Note that (A-1), (A-2), ..., (p-1) imply that both Eqs. (1) and (5) have unique continuous nonanticipating solutions for each $u(\cdot) \in \mathcal{U}$ and $E \int_0^\tau \|X_t\|^2 dt < \infty$, $E \int_0^\tau \|\phi_t\|^2 dt < \infty$. This follows directly from the existence theorems in [8] and [10].

THEOREM 1 (Stochastic Maximum Principle). *Assume the hypotheses (A-1), (A-2), ..., (p-1). Let $u(\cdot) \in \mathcal{U}$ and X_t be the corresponding trajectory (i.e., solution of (1)) and ϕ_t be the corresponding Lagrange multiplier (i.e., solution of (5)). Then in order for $u(\cdot)$ to minimize Φ in Eq. (3) over \mathcal{U} it is necessary that for almost all t in $[0, \tau]$,*

$$\inf_{u \in \mathcal{U}} E[\mathcal{H}(\phi_t, X_t, u; t)] = E[\mathcal{H}(\phi_t, X_t, u(t); t)]. \quad (7)$$

We now consider the existence problem. In this case we suppose G is a Hilbert space with inner product $\langle \cdot, \cdot \rangle_0$ and U a closed subspace. Then \mathcal{U} is also a Hilbert space with inner product $\langle u(\cdot), v(\cdot) \rangle_0 = \int_0^\tau \langle u(t), v(t) \rangle_0 dt$. \mathcal{U} is obviously separable. Let $k(s, t)$ be a symmetric square integrable kernel function in $[0, \tau] \times [0, \tau]$, i.e., $\int_0^\tau \int_0^\tau |k(s, t)|^2 ds dt < \infty$. Define an integral operator T from \mathcal{U} into itself by

$$Tu(s) = \int_0^\tau k(s, t) u(t) dt, \quad 0 \leq s \leq \tau. \quad (8)$$

Let $\mathcal{U}_{\tau, r}$ ($r > 0$) consist of all control functions of the form Tu , where $\|u\|_0 \leq r$.

THEOREM 2 (Existence for Optimal Controls). *In (4) suppose: (i) A satisfies (A-1) and (A-2), (ii) C and F are bounded continuous maps from $[0, \tau]$ into $L(B, H)$, and $D \in L(B, H)$, (iii) θ and ζ are bounded continuous maps from $[0, \tau] \times U$ into H , and (iv) g is a real-valued bounded continuous function on $[0, \tau] \times U$. Assume $DK(t, x) = 0$ for all t and x and $DC(t) = \alpha(t)D$ for some real-valued bounded continuous function α . Furthermore, suppose $X_0 = 0$ in Eq. (1). Then the functional Φ in Eq. (3) has both maximum and minimum over $\mathcal{U}_{\tau, r}$.*

Remark. $X_0 = 0$ is only a technical assumption for the proof.

3. PROOF OF THEOREM 1

We state a lemma which can be proved in the same way as Lemma A.2 [10].

LEMMA 1. *Suppose X_t is the solution of Eq. (1) corresponding to $u(\cdot) \in \mathcal{U}$. Then, for $0 \leq t, s \leq \tau$,*

$$E \|X_s - X_t\|^2 \leq c_1(e^{c_1|s-t|} - 1)(1 + E \|X_t\|^2 + \|u\|_0^2),$$

where c_1 is a constant depending only on the Lipschitzian constant c in the hypothesis and the norm $\|J\|_{B,B}$ of J .

The next lemma deals with the variation of controls. Let $u(\cdot) \in \mathcal{U}$ and X_t be the corresponding trajectory. Let t_0 be a Lebesgue point of $u(\cdot)$ (see, e.g., [7]) and $\epsilon > 0$, $z \in U$, define the perturbed control function $u_{\epsilon,z}(\cdot)$ of $u(\cdot)$ as follows:

$$u_{\epsilon,z}(t) = \begin{cases} u(t) & 0 \leq t \leq t_0 - \epsilon \\ z & t_0 - \epsilon < t \leq t_0 \\ u(t) & t_0 < t \leq \tau. \end{cases}$$

Note that $u_{\epsilon,z}(\cdot) \in \mathcal{U}$ and let $X_t^{\epsilon,z}$ be the corresponding trajectory such that $X_t^{\epsilon,z} = X_0$. We point out that $X_t^{\epsilon,z} - X_t$ is a stochastic process in H (rather than B). This will become clear through the following lemmas and the proofs.

LEMMA 2. Let t_0 be a Lebesgue point of $u(\cdot)$ then

$$E | X_{t_0}^{\epsilon,z} - X_{t_0} - \epsilon[\sigma(t_0, X_{t_0}, z) - \sigma(t_0, X_{t_0}, u(t_0))]|^2 = o(\epsilon^2).$$

Proof. By the uniqueness of solution it follows that $X_{t_0-\epsilon}^{\epsilon,z} = X_{t_0-\epsilon}$ since $X_0^{\epsilon,z} = X_0$. In order to simplify notation, set $Y_t = X_t^{\epsilon,z}$, $t_0 - \epsilon < t \leq t_0$. Then

$$Y_t = X_{t_0-\epsilon} + \int_{t_0-\epsilon}^t A(s, Y_s) dW_s + \int_{t_0-\epsilon}^t \sigma(s, Y_s, z) ds$$

and

$$X_t = X_{t_0-\epsilon} + \int_{t_0-\epsilon}^t A(s, X_s) dW_s + \int_{t_0-\epsilon}^t \sigma(s, X_s, u(s)) ds.$$

Let

$$\lambda(t) = Y_t - X_t - (t - t_0 + \epsilon)[\sigma(t_0, X_{t_0}, z) - \sigma(t_0, X_{t_0}, u(t_0))]. \quad (10)$$

It can be checked easily that

$$\lambda(t) = \text{I} + \text{II} - \text{III}, \quad (11)$$

where

$$\text{I} = \int_{t_0-\epsilon}^t [A(s, Y_s) - A(s, X_s)] dW_s,$$

$$\text{II} = \int_{t_0-\epsilon}^t [\sigma(s, Y_s, z) - \sigma(t_0, X_{t_0}, z)] ds,$$

and

$$\text{III} = \int_{t_0-\epsilon}^t [\sigma(s, X_s, u(s)) - \sigma(t_0, X_{t_0}, u(t_0))] ds.$$

In the following let c_2 denote a general constant depending only on c (Lipschitzian constant in the hypotheses) and c_1 (in Lemma 1).

Clearly,

$$\text{I} = \int_{t_0-\epsilon}^t [K(s, Y_s) - K(s, X_s)] dW_s,$$

and by Proposition 3.1 in [8],

$$\begin{aligned} E |\text{I}|^2 &= E \int_{t_0-\epsilon}^t \|K(s, Y_s) - K(s, X_s)\|_2^2 ds. \\ &\leq c_2 E \int_{t_0-\epsilon}^t \|Y_s - X_s\|^2 ds. \end{aligned}$$

Let

$$\beta(t) = Y_t - X_t \quad t_0 - \epsilon < t \leq t_0 \quad (12)$$

then

$$E |\text{I}|^2 \leq c_2 \int_{t_0-\epsilon}^t E \|\beta(s)\|^2 ds. \quad (13)$$

Writing $\sigma(s, Y_s, z) - \sigma(t_0, X_{t_0}, z)$ as $[\sigma(s, Y_s, z) - \sigma(s, X_{t_0}, z)] + [\sigma(s, X_{t_0}, z) - \sigma(t_0, X_{t_0}, z)]$ and using $(\sigma-1)$, we can estimate II as follows:

$$E |\text{II}|^2 \leq h(t) + c_2 \int_{t_0-\epsilon}^t E \|\beta(s)\|^2 ds, \quad (14)$$

where

$$h(t) = c_2 \epsilon \int_{t_0-\epsilon}^t [E \|X_s - X_{t_0}\|^2 + E |\sigma(s, X_{t_0}, z) - \sigma(t_0, X_{t_0}, z)|^2] ds.$$

Similarly, we have

$$E |\text{III}|^3 \leq k(t) + c_2 \int_{t_0-\epsilon}^t E \|\beta(s)\|^2 ds, \quad (15)$$

where

$$\begin{aligned} k(t) &= c_2 \epsilon \int_{t_0-\epsilon}^t [E \|X_s - X_{t_0}\|^2 + E |\sigma(s, X_{t_0}, u(t_0)) - \sigma(t_0, X_{t_0}, u(t_0))|^2 \\ &\quad + |u(s) - u(t_0)|_0^2] ds. \end{aligned}$$

Putting (11), (13), (14), and (15) together, we obtain

$$E \|\lambda(t)\|^2 \leq h(t) + k(t) + c_2 \int_{t_0-\epsilon}^t E \|\beta(s)\|^2 ds. \quad (16)$$

On the other hand, from (1) we get

$$Y_t - X_t = \lambda(t) + (t - t_0 + \epsilon)[\sigma(t_0, X_{t_0}, z) - \sigma(t_0, X_{t_0}, u(t_0))],$$

whence

$$\begin{aligned} E \|\beta(t)\|^2 &\leq c_2 E \|\lambda(t)\|^2 + c_2 \epsilon^2 E \|\sigma(t_0, X_{t_0}, z) - \sigma(t_0, X_{t_0}, u(t_0))\|^2 \\ &\leq c_2 E \|\lambda(t)\|^2 + c_2 \epsilon^2 E \|\sigma(t_0, X_{t_0}, z) - \sigma(t_0, X_{t_0}, u(t_0))\|^2 \\ &\leq c_2 \int_{t_0-\epsilon}^t E \|\beta(s)\|^2 ds + h(t) + k(t) + c_2 \epsilon^2 \|z - u(t_0)\|_0^2 \\ &\quad \text{by (16) and } (\sigma-1). \end{aligned}$$

Thus by Gronwall's inequality,

$$\begin{aligned} E \|\beta(t)\|^2 &\leq h(t) + k(t) + c_2 \epsilon^2 \|z - u(t_0)\|_0^2 \\ &\quad + c_2 \int_{t_0-\epsilon}^t e^{c_2(t-s)} [h(s) + k(s) + c_2 \epsilon^2 \|z - u(t_0)\|_0^2] ds. \end{aligned}$$

Note that $h(t)$ and $k(t)$ are increasing functions of t , hence from the above inequality we have

$$E \|\beta(t)\|^2 \leq c_2 [h(t_0) + k(t_0) + \epsilon^2 \|z - u(t_0)\|_0^2]. \quad (17)$$

From (16) and (17) it is easy to see that

$$E \|\lambda(t_0)\|^2 \leq c_2 [h(t_0) + k(t_0) + \epsilon^3 \|z - u(t_0)\|_0^2].$$

To finish the proof, simply note that both $h(t_0)$ and $k(t_0)$ are $o(\epsilon^2)$ by Lemma 1 and the fact that $E \|\sigma(s, X_{t_0}, z) - \sigma(t_0, X_{t_0}, z)\|^2$ and $E \|\sigma(s, X_{t_0}, u(t_0)) - \sigma(t_0, X_{t_0}, u(t_0))\|^2$ are continuous functions of s , and also, note that $\int_{t_0-\epsilon}^{t_0} \|u(s) - u(t_0)\|_0^2 ds = o(\epsilon)$ since t_0 is a Lebesgue point of $u(\cdot)$.

LEMMA 3. If t_0 is a Lebesgue point of $u(\cdot)$ and $E[f(\cdot, X_{\cdot}, u(\cdot))]$ then

$$\begin{aligned} \lim_{\epsilon \downarrow 0} \epsilon^{-1} E \int_{t_0-\epsilon}^{t_0} [f(t, X_t^{\epsilon, z}, z) - f(t, X_t, u(t))] dt \\ = E[f(t_0, X_{t_0}, z) - f(t_0, X_{t_0}, u(t_0))]. \end{aligned}$$

Proof. Let

$$\xi(t) = E[f(t, X_t, u(t))], \quad 0 \leq t \leq \tau.$$

It follows from (f-1) that

$$|\xi(t)| \leq c(1 + E\|X_t\|^2 + |u(t)|_0^2).$$

Lemma 1 gives $\int_0^\tau E\|X_t\|^2 dt < \infty$, hence $\int_0^\tau |\xi(t)| dt < \infty$. Therefore,

$$\lim_{\epsilon \downarrow 0} \epsilon^{-1} E \int_{t_0-\epsilon}^{t_0} f(t, X_t, u(t)) dt = E[f(t_0, X_{t_0}, u(t_0))]. \quad (18)$$

Similarly,

$$\lim_{\epsilon \downarrow 0} \epsilon^{-1} E \int_{t_0-\epsilon}^{t_0} f(t, X_t, z) dt = E[f(t_0, X_{t_0}, z)]. \quad (19)$$

On the other hand, since f is C_H^1 in x variable,

$$f(t, X_t^{\epsilon, z}, z) - f(t, X_t, z) = \langle f_x(t, X_t, z), X_t^{\epsilon, z} - X_t \rangle + o(\|X_t^{\epsilon, z} - X_t\|).$$

Use (f-2) and apply Lemma 2 to conclude that

$$\lim_{\epsilon \downarrow 0} \epsilon^{-1} \int_{t_0-\epsilon}^{t_0} E |f(t, X_t^{\epsilon, z}, z) - f(t, X_t, z)|^2 dt = 0. \quad (20)$$

Equations (18), (19), and (20) give the assertion of the lemma.

In the interval $t_0 < t \leq \tau$, X_t and $X_t^{\epsilon, z}$ satisfy the same stochastic differential equation (with initial distributions X_{t_0} and $X_{t_0}^{\epsilon, z}$ at t_0 , respectively), namely,

$$dV_t = A(t, V_t) dW_t + \sigma(t, V_t, u(t)) dt. \quad (21)$$

Now we will employ some results in [10] to study Eq. (21). Let us recall the notion of mean-square differentiability in H -directions (MS - H -differentiability) introduced in [10, Definition 5]. With a slight modification of Theorem 7 [10] in order to apply to our case here we conclude that the solution of Eq. (21) is MS - H -differentiable. The MS - H -derivative of V_t at X_{t_0} is given by the solution of the following $L(H, H)$ -valued stochastic equation, $t_0 \leq t \leq \tau$,

$$Y_{t_0, t} = I + \int_{t_0}^t Y_{t_0, s} \Delta K_x(s, X_s) dW_s + \int_{t_0}^t Y_{t_0, s} \circ \sigma_x(s, X_s, u(s)) ds, \quad (22)$$

where I is the identity operator of H .

Remark. For $S \in L(H, L(H, H))$ and $T \in L(H, H)$, $T \Delta S \in L(H, L(H, H))$ is defined by: $T \Delta S(h) = T \circ [S(h)]$, $h \in H$, where \circ is the composition of two operators. We define also that $S \Delta T(h) = S(h) \circ T$. For a detailed discussion of the above operator-valued stochastic integral equation, see [9 and 10].

LEMMA 4. Let t_0 be a Lebesgue point of $u(\cdot)$ then

$$\begin{aligned} & \lim_{\epsilon \downarrow 0} \epsilon^{-1} E \int_{t_0}^{\tau} [f(t, X_t^{\epsilon, z}, u(t)) - f(t, X_t, u(t))] dt \\ &= E \int_{t_0}^{\tau} \langle f_x(t, X_t, u(t)), Y_{t_0, t}(\sigma(t_0, X_{t_0}, z) - \sigma(t_0, X_{t_0}, u(t_0))) \rangle dt, \end{aligned}$$

where $Y_{t_0, t}$ is the solution of Eq. (22).

Proof. First note that using the same argument in the proof of Lemma 2 we see that $E |X_t^{\epsilon, z} - X_t|^2$ is dominated by $E |X_{t_0}^{\epsilon, z} - X_{t_0}|^2$ which is $o(\epsilon^2)$ by Lemma 2. Hence $E |X_t^{\epsilon, z} - X_t|$ is $o(\epsilon)$ uniformly for $t_0 < t \leq \tau$. Therefore,

$$\begin{aligned} & \lim_{\epsilon \downarrow 0} \epsilon^{-1} E \int_{t_0}^{\tau} [f(t, X_t^{\epsilon, z}, u(t)) - f(t, X_t, u(t))] dt \\ &= \lim_{\epsilon \downarrow 0} \epsilon^{-1} E \int_{t_0}^{\tau} \langle f_x(t, X_t, u(t)), X_t^{\epsilon, z} - X_t \rangle dt. \end{aligned} \quad (23)$$

Let $Y_{t_0, t, x}$ denote the MS-H-derivative of V_t at x then

$$X_t^{\epsilon, z} - X_t = \int_0^1 Y_{t_0, t, X_{t_0} + \lambda(X_{t_0}^{\epsilon, z} - X_{t_0})}(X_{t_0}^{\epsilon, z} - X_{t_0}) d\lambda, \quad t_0 < t \leq \tau.$$

Using (A-3) and (σ -2) to show

$$E \| Y_{t_0, t, X_{t_0} + \lambda(X_{t_0}^{\epsilon, z} - X_{t_0})} \|_{H, H}$$

is uniformly bounded for $\epsilon > 0$ and then Lebesgue dominated convergence theorem and Lemma 2 to find the limit, we conclude that almost surely

$$\lim_{\epsilon \downarrow 0} \epsilon^{-1} (X_t^{\epsilon, z} - X_t) = Y_{t_0, t, X_{t_0}}(\sigma(t_0, X_{t_0}, z) - \sigma(t_0, X_{t_0}, u(t_0))),$$

which together with (23) yield the lemma. Note that $Y_{t_0, t, X_{t_0}}$ is the solution of Eq. (22).

LEMMA 5.

$$\begin{aligned} \lim_{\epsilon \downarrow 0} \epsilon^{-1} E[p(X_\tau^{\epsilon, z}) - p(X_\tau)] \\ = E\langle p'(X_\tau), Y_{t_0, \tau}(\sigma(t_0, X_{t_0}, z) - \sigma(t_0, X_{t_0}, u(t_0))) \rangle, \end{aligned}$$

where $Y_{t_0, t}$ is the same as Lemma 4, i.e., the solution of Eq. (22).

Proof. Similar to the previous one.

From now on, we assume $u(\cdot)$ is an optimal control minimizing Φ in Eq. (3) and X_t the corresponding optimal trajectory. Define

$$\Lambda_{z, t_0}(\epsilon) = \begin{cases} \Phi(u_{\epsilon, z}, X_{\epsilon, z}), & \epsilon > 0 \\ \Phi(u, X), & \epsilon = 0 \end{cases} \quad (24)$$

where $z \in U$ and $u_{\epsilon, z}$ is given in (9) and $X_{\epsilon, z}$ is the corresponding trajectory with $X_0^{\epsilon, z} = X_0$.

PROPOSITION 1. *If t_0 is a Lebesgue point of $u(\cdot)$ and $E[f(\cdot, X_\cdot, u(\cdot))]$ then the right derivative $\Lambda'_{z, t_0}(0)$ of Λ_{z, t_0} at the origin exists. Furthermore, it is given by:*

$$\begin{aligned} \Lambda'_{z, t_0}(0) &= E[f(t_0, X_{t_0}, z) - f(t_0, X_{t_0}, u(t_0))] \\ &+ E \int_{t_0}^{\tau} \langle f_x(t, X_t, u(t)), Y_{t_0, t}(\sigma(t_0, X_{t_0}, z) - \sigma(t_0, X_{t_0}, u(t_0))) \rangle dt \\ &+ E\langle p'(X_\tau), Y_{t_0, \tau}(\sigma(t_0, X_{t_0}, z) - \sigma(t_0, X_{t_0}, u(t_0))) \rangle, \end{aligned} \quad (25)$$

where $Y_{t_0, t}(t_0 \leq t \leq \tau)$ is the solution of Eq. (22).

Proof. Note that in the interval $0 \leq t \leq t_0 - \epsilon$, $u_{\epsilon, z}(t) = u(t)$. Hence by the uniqueness, $X_t^{\epsilon, z} = X_t$. Hence

$$E \int_0^{t_0 - \epsilon} [f(t, X_t^{\epsilon, z}, u_{\epsilon, z}(t)) - f(t, X_t, u(t))] dt = 0.$$

This rest follows immediately from Lemmas 3, 4, and 5.

Define a stochastic process with state space H by:

$$\phi_s = \int_s^\tau Y_{s, t}^*(f_x(t, X_t, u(t))) dt + Y_{s, \tau}^*(p'(X_\tau)), \quad 0 \leq s \leq \tau. \quad (26)$$

Use this ϕ_s and the Hamiltonian \mathcal{H} defined in (6), (25) can then be rewritten as,

$$\Lambda'_{z, t_0}(0) = E\mathcal{H}(\phi_{t_0}, X_{t_0}, z; t_0) - E\mathcal{H}(\phi_{t_0}, X_{t_0}, u(t_0); t_0).$$

Now, in order for $u(\cdot)$ to minimize Φ , it is necessary to have $A'_{z,t_0}(0) \geq 0$, i.e.,

$$E\mathcal{H}(\phi_{t_0}, X_{t_0}, z; t_0) \geq E\mathcal{H}(\phi_{t_0}, X_{t_0}, u(t_0); t_0)$$

for all $z \in U$ and all the Lebesgue points t_0 of $u(\cdot)$ and $E[f(\cdot, X, u(\cdot))]$. Clearly, for all those t_0 ,

$$\inf_{z \in U} E\mathcal{H}(\phi_{t_0}, X_{t_0}, z; t_0) = E\mathcal{H}(\phi_{t_0}, X_{t_0}, u(t_0); t_0).$$

Of course, it is well-known that those t_0 's are almost everywhere in $[0, \tau]$. Thus we will finish the proof if we show that ϕ_s defined in (26) satisfies Eq. (5). First note that from Eq. (22) $Y_{\tau,\tau} = I$, the identity operator of H . Hence $\phi_\tau = p'(X_\tau)$. From (26), we derive easily the stochastic differential of ϕ_s ,

$$\begin{aligned} d\phi_s = & -Y_{s,s}^*(f_x(s, X_s, u(s))) ds + \int_s^\tau (d_s Y_{s,t}^*)(f_x(t, X_t, u(t))) dt \\ & + (d_s Y_{s,\tau}^*)(p'(X_\tau)), \end{aligned} \quad (27)$$

where d_s indicates the differentials with respect to s -variable. On the other hand, from Eq. (22) we have

$$Y_{s,t} = I + \int_s^t Y_{s,\lambda} \Delta K_x(\lambda, X_\lambda) dW_\lambda + \int_s^t Y_{s,\lambda} \circ \sigma_x(\lambda, X_\lambda, u(\lambda)) ds,$$

and, taking adjoint, we have

$$Y_{s,t}^* = I + \int_s^t K_x(\lambda, X_\lambda)^* \Delta Y_{s,\lambda}^* dW_\lambda + \int_s^t \sigma_x(\lambda, X_\lambda, u(\lambda))^* \circ Y_{s,\lambda}^* d\lambda,$$

where if $S \in L(H, L(H, H))$ then $S^* \in L(H, L(H, H))$ is defined to be $S^*(h) = [S(h)]^*$, $h \in H$. The above equation can be written in the stochastic differential form, $s \leq t \leq \tau$,

$$d_s Y_{s,t}^* = -K_x(s, X_s)^* \Delta Y_{s,t}^* dW_s - \sigma_x(s, X_s, u(s))^* \circ Y_{s,t}^* ds. \quad (28)$$

Substitute (28) into (27) and note that $Y_{s,s} = I$, it follows that

$$d\phi_s = -f_x(s, X_s, u(s)) ds - K_x(s, X_s)^* \phi_s dW_s - \sigma_x(s, X_s, u(s))^* \phi_s ds.$$

Hence ϕ_s satisfies Eq. (5). (Note that $A_x = K_x$.)

4. EXAMPLE

EXAMPLE 1. Consider the system

$$dX_t = dW_t - (CX_t + u(t)z_0) dt, \quad 0 \leq t \leq \tau, \quad (29)$$

where $C \in L(B, H)$ and $z_0 \in H$. The control region is $[-1, 1]$ in the real line. The cost function is given by $f(t, x, u) = \langle Ax, z_0 \rangle + u^2$ and $p \equiv 0$. In this case, the adjoint system is,

$$d\phi_t = (C^*\phi_t - A^*z_0) dt, \quad \phi_\tau = 0.$$

The solution is easily seen to be

$$\phi_t = (\tau - t) e_*((t - \tau) C^*) A^* z_0, \quad (30)$$

where $e_*(S) = \sum_{n=0}^{\infty} S^n / (n + 1)!$ for $S \in L(H, H)$.

Let $c_1(t) = \langle \phi_t, z_0 \rangle$ and $c_2(t) = E\langle AX_t, z_0 \rangle - E\langle \phi_t, CX_t \rangle$, then we have

$$E\mathcal{H}(\phi_t, X_t, u; t) = u^2 - c_1(t)u + c_2(t).$$

Obviously,

$$\inf_{-1 \leq u \leq 1} E\mathcal{H}(\phi_t, X_t, u; t) = \begin{cases} -(c_1(t)^2/4) + c_2(t), & \text{if } |c_1(t)| \leq 2; \\ 1 - c_1(t) + c_2(t), & \text{if } c_1(t) > 2; \\ 1 + c_1(t) + c_2(t), & \text{if } c_1(t) < -2. \end{cases}$$

Therefore by Theorem 1, in order for $u(t)$ to be optimal it must be

$$u(t) = \begin{cases} c_1(t)/2, & \text{if } |c_1(t)| \leq 2; \\ 1, & \text{if } c_1(t) > 2; \\ -1, & \text{if } c_1(t) < -2. \end{cases}$$

The corresponding trajectory, i.e., the solution of (29), is given by

$$X_t = e^{-Ct}x + \int_0^t e^{C(s-t)} dW_s - \left[\int_0^t e^{C(s-t)} u(s) ds \right] z_0.$$

As a matter of fact, by putting $u(t)$ and X_t into Φ in (3) it can be checked that $u(\cdot)$ is the optimal control.

EXAMPLE 2. Consider one dimensional case, i.e., $B = H = \mathbb{R}$. Let b_t be one dimensional Brownian motion starting at the origin. Consider the system with $x_0 = 0$,

$$dx_t = x_t db_t + \{(1/2)x_t + u(t)\} dt, \quad 0 \leq t \leq \tau.$$

The control region U is \mathbb{R} and the cost function is quadratic, i.e., $f(t, x, u) = x^2 + xu + u^2$, $p = 0$.

Take expectation in both sides of the system, it follows immediately that

$$Ex_t = \int_0^t e^{(t-s)/2} u(s) ds. \quad (31)$$

The adjoint system is

$$\begin{aligned} d\phi_t &= -\phi_t db_t - [(1/2)\phi_t + 2x_t + u(t)] dt, \\ \phi_\tau &= 0. \end{aligned}$$

We have

$$E\phi_t = \int_t^\tau e^{(s-t)/2} [2Ex_s + u(s)] ds. \quad (32)$$

Now, the Hamiltonian is

$$\mathcal{H}(\phi_t, x_t, u; t) = u^2 + (x_t + \phi_t)u + x_t^2 + (1/2)x_t\phi_t.$$

Hence

$$\inf_{-\infty < u < \infty} E\mathcal{H}(\phi_t, x_t, u; t) = \beta(t) - \alpha(t)^2/4,$$

where $\alpha(t) = Ex_t + E\phi_t$ and $\beta(t) = Ex_t^2 + (1/2)Ex_t\phi_t$.

Therefore, by Theorem 1 in order for $u(\cdot)$ to be optimal we must have

$$u(t) = -\alpha(t)/2.$$

From the above equation, (31) and (32) we can derive a differential equation for $u(t)$, i.e.,

$$u''(t) = 3/4 u(t)$$

with "initial" conditions

$$u(0) + 2u'(0) = 0 \quad \text{and} \quad 3u(\tau) + 2u'(\tau) = 0.$$

It is then easy to see that $u \equiv 0$. Obviously, this $u(\cdot)$ is the optimal control by noting that $\Phi \geq 0$.

5. PROOF OF THEOREM 2

LEMMA 6. $EX_t = \int_0^t [\exp(\int_s^t C(\lambda) d\lambda)] \theta(s, u(s)) ds$, where $C(\lambda)$ is regarded as an operator of H .

Remark. Although X_t is a process in B , EX_t is a process in H .

Proof. From the following equation

$$X_t = \int_0^t A(s, X_s) dW_s + \int_0^t [C(s) X_s + \theta(s, u(s))] ds,$$

we have, since $E \int_0^t A(s, X_s) dW_s = 0$ (see [8]),

$$EX_t = \int_0^t [C(s) EX_s + \theta(s, u(s))] ds.$$

The lemma follows by solving the above H -valued integral equation.

LEMMA 7. $E |DX_t|^2 = \int_0^t [\exp(2 \int_s^t \alpha(\lambda) d\lambda)] h(s) ds$, where $h(s) = 2\langle EX_s, D^*D\theta(s, u(s)) \rangle + \text{trace } D^*DJJ^*$.

Proof. Let $\phi(x) = |Dx|^2$. By Ito's formula [8],

$$\begin{aligned} d\phi(X_t) &= (A^*(t, X_t) \phi'(X_t), dW_t) + \{\langle \phi'(X_t), C(t) X_t + \theta(t, u(t)) \rangle \\ &\quad + (1/2) \text{trace}[A^*(t, X_t) \phi''(X_t) A(t, X_t)]\} dt. \end{aligned} \quad (33)$$

But $\phi'(x) = 2D^*Dx$ and $\phi''(x) = 2D^*D$. Therefore,

$$\begin{aligned} \langle \phi'(X_t), C(t) X_t \rangle &= 2\langle D^*DX_t, C(t) X_t \rangle, \\ &= 2\langle DX_t, DC(t) X_t \rangle, \\ &= 2\langle DX_t, \alpha(t) DX_t \rangle, \\ &= 2\alpha(t) |DX_t|^2, \\ &= 2\alpha(t) \phi(X_t), \end{aligned}$$

and

$$(1/2) \text{trace}[A^*(t, X_t) \phi''(X_t) A(t, X_t)] = \text{trace } D^*DJJ^*.$$

Here we have used the assumption that $DC(t) = \alpha(t)D$ and $DK(t, x) = 0$. Note that $E \int_0^t (A^*(s, X_s) \phi'(X_s), dW_s) = 0$ (see [8]). Hence from Eq. (33) we have

$$\begin{aligned} E\phi(X_t) &= 2 \int_0^t \alpha(s) E\phi(X_s) ds \\ &\quad + \int_0^t \{2\langle EX_s, D^*D\theta(s, u(s)) \rangle + \text{trace } D^*DJJ^*\} ds. \end{aligned}$$

The lemma then follows easily by solving the above ordinary integral equation.

Now, from Lemma 6 and Lemma 7 it follows that the functional Φ in (3), defined in the function space \mathcal{U} , is continuous. This can be checked as follows: If $u_n(\cdot)$ converges to $u(\cdot)$ in \mathcal{U} then $u_n(\cdot)$ converges to $u(\cdot)$ almost everywhere. But every map is bounded and continuous, therefore Lebesgue dominated convergence theorem can be applied to conclude that $\Phi(u_n)$ converges to $\Phi(u)$.

Finally, we recall some facts about abstract Wiener space. \mathcal{U} is a real separable Hilbert space and the operator T defined in (8) is a Hilbert-Schmidt operator of \mathcal{U} . Let $\hat{\mathcal{U}} = (\ker T)^\perp$. Then T is a one-to-one Hilbert-Schmidt operator of $\hat{\mathcal{U}}$. It is well-known that $(T(\hat{\mathcal{U}}), \hat{\mathcal{U}})$ is an abstract Wiener space and that the set $\{Tu; u \in \hat{\mathcal{U}}, \|u\|_0 \leq r\}$ is compact in $\hat{\mathcal{U}}$ -topology. Clearly this set is also compact in \mathcal{U} . But this set is the same as $\{Tu; u \in \mathcal{U}, \|u\|_0 \leq r\} = \mathcal{U}_{T,r}$. Therefore, we conclude that $\mathcal{U}_{T,r}$ is compact in \mathcal{U} . Now, since Φ is continuous in \mathcal{U} , Φ assumes both maximum and minimum over $\mathcal{U}_{T,r}$.

REFERENCES

1. A. V. BALAKRISHNAN, Stochastic control: a function space approach, *SIAM J. Control* 10 (1972), 285-297.
2. W. H. FLEMING, Stochastic Lagrange multipliers, in "Mathematical theory of control" A. V. Balakrishnan and L. W. Neustadt, (Eds.), Academic Press, New York, 1967.
3. W. H. FLEMING, Optimal control of partially observable diffusions, *SIAM J. Control* 6 (1968), 194-214.
4. W. H. FLEMING, Optimal continuous-parameter stochastic control, *SIAM Review* 11 (1969), 470-509.
5. W. H. FLEMING AND M. NISIO, On the existence of optimal stochastic controls, *J. Math. Mech.* 15 (1966), 777-794.
6. L. GROSS, Abstract Wiener spaces, *Proc. 5th Berkeley Sym. Math. Stat. Prob.* (1965), 31-42.
7. E. HEWITT AND K. STROMBERG, "Real and Abstract Analysis," Springer-Verlag, New York, 1965.
8. H.-H. KUO, Stochastic integrals in abstract Wiener space, *Pacific J. Math.* 41 (1972), 469-483.
9. H.-H. KUO, On operator-valued stochastic integrals, *Bull. Amer. Math. Soc.* 79 (1973), 207-210.
10. H.-H. KUO, Stochastic integrals in abstract Wiener space II: Regularity properties, *Nagoya Math. J.* (to appear).
11. H. J. KUSHNER, On the stochastic maximum principle: Fixed time of control, *J. Math. Anal. Appl.* 11 (1965), 78-92.

12. L. S. PONTRYAGIN, V. G. BOLTYANSKII, R. V. GAMKREJIDZE, AND E. F. MISCHENKO, "The Mathematical Theory of Optimal Processes" (trans. Trirogoff K.N.), Interscience Publishers, New York, 1962.
13. V. M. WARFIELD, "A Stochastic Maximum Principle," Ph.D. thesis, Brown University, 1971.